


# Differences between the values of frequencies by different fitters

## 16CygA&B and Kepler Legacy values

Ian Roxburgh, Queen Mary University of London

The differences between the oscillation frequencies and uncertainty estimates of a star derived by different fitters can be large, sufficiently large so that, were one to find a stellar model that fitted one frequency set ( $\chi^2 \sim 1$ ), it does not fit an alternative set. The table below gives 21 examples, comparing frequency sets in common between the Kepler Legacy project and frequency sets from Appourchaux et al (2014)<sup>1</sup> and Davies et al (2015)<sup>2</sup>. Figure 1 displays the frequency differences  $\nu_L - \nu_D$  (Legacy-Davies) for 16CygA&B and the  $\chi^2$  of the fits to each other. A model whose frequencies fit the Legacy frequency set for 16CygA with  $\chi_L^2 < 1$  could have  $\chi_D^2 > 10$  for a fit to Davies's frequency set and so would be rejected.



These differences are not statistical uncertainties; given the same input light curve, differences in estimated frequencies are due to different assumptions/constraints in mode fitting techniques, the segment of the time series used, and the algorithms for determining power spectra. The differences constitute uncertainties in the values of the frequencies and should be added to estimates of errors.

To better understand these differences I applied my own mode fitting code (described below) to 16CygA&B and KIC 6116408, 8379927 and 10454113, using the kasoc power spectra, Davies's spectra and my own power spectra derived from kasoc light curves. I find significant differences between frequencies derived from different power spectra and a smaller difference between weighted and unweighted spectra. I find that different mode height ratios  $h_l/h_0$  (fixed or free) have little effect except for low values; that too low an inclination angle can have a significant effect; and that rejecting modes with low signal to noise gives very much better agreement between different determinations of frequencies.. Details are presented below.

For 16CygA&B I find modest agreement between Davies's ( $\nu_D$ ) and my frequencies ( $\nu_{RD}$ ) using Davies's power spectra ( $\chi_A^2 = 0.33, \chi_B^2 = 0.21$ ) and very good agreement if I reject modes with low signal/noise ( $\chi_A^2 = 0.06, \chi_B^2 = 0.03$ ). I do not find such agreement between the Legacy frequencies ( $\nu_L$ ) and my frequencies ( $\nu_{RL}$ ) derived from the Legacy power spectrum for 16CygA ( $\chi_A^2 = 1.53$ ). I find much better agreement between my values  $\nu_{RD}, \nu_{RL}$  from the two power spectra ( $\chi^2 = 0.44$ ). I show that there are some misfits in the Legacy frequencies for 16CygA (Figure 4 below).

For 16CygB two versions of the power spectrum (v1,v2) have been listed on the KASOC website, the earlier version (v1) gives modest agreement between my  $\nu_{RL}$  and the legacy values ( $\chi_B^2 = 0.35$ ) but these differ substantially from the values  $\nu_{RD}$  from Davies's spectrum ( $\chi^2 = 1.04$ ); using (v2) gives values closer to Davies's values ( $\chi^2 = 0.27$ ) but a considerably larger difference from the Legacy values  $\nu_L$  ( $\chi^2 = 1.13$ ).


For the other 3 stars I find closer (but not good) agreement between my frequencies ( $\nu_{RA}$ ) and Appourchaux's ( $\nu_A$ ) than with the Legacy values ( $\nu_L$ ).

## $\chi^2$ of fits of Legacy frequencies to those of Appourchaux and Davies

$\chi_m^2$  is the fit of Legacy frequencies  $\nu_L \pm \sigma_L$  to Davies's  $\nu_D \pm \sigma_D$ , or Appourchaux's  $\nu_A \pm \sigma_A$ , taking  $\sigma_m = \max(\sigma_L, \sigma_D)$  or  $\max(\sigma_L, \sigma_A)$ ,  $\chi_L^2$  with  $\sigma_L$ ,  $\chi_D^2$  with  $\sigma_D$ ,  $\chi_A^2$  with  $\sigma_A$ .  $n(\nu)$  is the number of frequencies in common,  $n(< \sigma_m)$  the number of frequencies that agree within  $1\sigma_m$ . 4 examples are shown below.

Fitter	KIC no	$\Delta$	$n(\nu)$	$n(< \sigma_m)$	$\chi_m^2$	$\chi_L^2$	$\chi_D^2 / \chi_A^2$
Davies	16CygA	103	50	23	1.60	1.64	11.47
Davies	16CygB	117	51	33	1.35	1.62	1.79
Davies	8379927	120	49	36	0.50	0.94	0.59
App	12317678	64	52	22	1.65	2.42	4.16
App	12258514	75	45	34	0.87	1.23	1.06
App	12009504	88	43	30	0.85	1.08	1.25
App	11081729	90	40	25	2.87	5.40	8.50
App	10454113	105	49	34	1.05	1.31	1.67
App	10162436	56	48	26	1.53	1.83	1.88
App	9812850	65	48	31	1.55	1.92	3.11
App	9206432	85	49	36	0.86	0.98	1.44
App	9139163	81	55	39	1.51	2.63	1.90
App	9139151	117	34	26	0.67	0.86	1.17
App	8694723	75	53	40	1.01	1.47	1.22
App	8379927	120	45	35	0.71	1.15	0.73
App	7206837	79	43	24	2.38	2.70	3.50
App	7103006	60	53	33	2.11	2.54	3.42
App	6679371	51	54	33	1.21	1.83	1.82
App	6508366	51	50	34	1.33	1.80	1.88
App	6116048	101	42	23	1.63	2.23	1.77
App	2837475	76	51	33	1.10	1.56	1.69
App	1435467	70	45	31	2.98	3.79	3.27

Frequency differences between Legacy ( $\nu_L \pm \sigma_L$ ) and Appourchaux ( $\nu_A \pm \sigma_A$ ) values









## iwr's frequencies ( $\nu_{RL}, \nu_{RD}$ ) vs Legacy ( $\nu_L$ ) and Davies ( $\nu_D$ ) values for 16CygA&B

**iwr's mode fitting algorithm.** For given rotational parameters ( $\omega, i$ ) I search for a global fit of symmetric Lorentzian profiles to a section of the power spectrum that includes the region of p-mode power, iteratively updating mode pairs to reach a minimum of a maximum likelihood estimator. The background is determined by fitting a Harvey type function to the high and low frequency extremes and to power minus mode power in windows in the central region and updated after each global iteration.. The starting values are given by local fitting of mode pairs. The mode heights and widths for  $\ell = 1, 2, 3$  are determined by interpolation in the values for  $\ell = 0$  with given (or free) mode height ratios. This procedure is then repeated with different ( $\omega, i$ ) to find the best fit. In fitting 16CygA&B I took Davies's central values for ( $\omega, i$ )=(0.496, 56) for A and (0.466, 36) for B, and constant mode height ratios  $h_1/h_0 = 1.554, h_2/h_0 = 0.582, h_3/h_0 = 0.040$ .


The figures show the differences between my frequencies  $\nu_{RD}, \nu_{RL}$  from the Davies and KASOC power spectra for 16CygA & B(v2). There is modest agreement with Davies's values for both 16CygA&B (except for modes with low signal/noise) but not with the Legacy values. Moreover my frequencies from the 2 power spectra  $\nu_{RD}, \nu_{RL}$  are in modest agreement with each other ( $\chi^2_A = 0.44, \chi^2_B = 0.27$ ). This suggests there could be some misfits in the Legacy mode fitting, which is clearly seen for 16CygA on the next page.

I also find a closer fit to Appourchaux's values for KIC6116084, 8379927, 10454113 than to Legacy values.



## 16CygA: iwr's fit to the Legacy power spectrum and Legacy frequencies and fit to Davies power spectrum and Davies frequencies

The following figure shows (in red) my fit to the full power spectrum used in the Legacy fit (courtesy of M Lund) [kplr012069424\_kasoc-wpsd\_slc\_v1.pow] overlaid on a  $0.2\mu\text{Hz}$  boxcar of the spectrum around 3 mode pairs and and (in blue) the location of the Legacy frequencies, which are not in agreement. Below is the comparable fit to the Davies power spectrum for 2 of the mode pairs, which are in agreement. This suggests there may be some error in the Legacy fitting algorithm.




## Frequencies at low signal to noise

As shown above I reproduce Davies's frequencies using Davies's power spectra for 16CygA&B with a  $\chi_A^2 = 0.33$  and  $\chi_B^2 = 0.21$ , the major divergences being at low and high frequencies where the mode heights are small compared to the background and so are very sensitive to modelling of the background, and to the derivation of power spectra from light curves. This, and large mode widths at high frequencies, makes me question the reliability of frequency estimates for low signal to noise.


I define signal/noise (S/N) as the maximum height of a (rotationally split) mode divided by the local background; this is shown in the left panels of the following figure for 16CygA&B for the fits to Davies's power spectra and 16CygB for the kasoc\_v2 power spectrum; all  $\ell = 3$  and some  $\ell = 0, 1, 2$  modes have  $S/N < 1$ ; as shown in the top 2 right panels if these modes are excluded the quality of the fit of my frequencies ( $\nu_{RD}$ ) to those of Davies ( $\nu_D$ ) is much improved ( $\chi_A^2 = 0.06$ ,  $\chi_B^2 = 0.03$ ).

The situation is different for 16CygB (and A not shown) using the kasoc\_v2 power spectrum - there is no improvement in the fit of my frequencies  $\nu_{RL}$  to the Legacy values ( $\nu_L$ ) - indeed the  $\chi^2$  of the fit it is slightly worse than when low S/N modes are included.

16CygA&B: iwr S/N=maximum of mode power/background (Davies pspec) and frequency differences



16CygB: iwr S/N=maximum mode power/background (kasoc\_v2 pspec) and frequency differences




## Dependence of frequencies on power spectra derived from kasoc light curves

The Legacy power spectra for 16CygA&B were derived from weighted kasoc light curves from Q6-Q17.2 whereas Davies used data from Q7-Q16 with smoothing. To explore the dependence of the frequencies on power spectra I derived spectra and frequencies  $\nu_{DT}$  from Davies's time series (private communication) and 3 sets from the kasoc light curves:  $\nu_{12}$  with data from Q6-Q17.2 weighted by the inverse of the flux error;  $\nu_{10}$  from Q7-Q16 also weighted, and  $\nu_n$  unweighted from Q7-Q16. I removed bad data (Inf and zero), had no gap filling, used the Lomb-Scargle algorithm and determined frequencies with the same routine as fits to the Legacy and Davies power spectra ( $\nu_{RD}, \nu_{RL}$ ). The top 2 panels show near perfect fits of  $\nu_{DT}$  to my  $\nu_{RD}$  for all frequencies indicating that there is no error in my power spectrum routine.

The bottom 4 panels show the best fits of frequencies from the my power spectra ( $\nu_{12}, \nu_{10}, \nu_n$ ) from the kasoc light curves to those from from Davies's and Legacy power spectra taking the Legacy uncertainties on frequencies  $\chi^2$  is for modes with  $S/N \geq 1$  and  $\chi_f^2$  for all frequencies. Not surprisingly the  $\nu_{RD}$  best fit  $\nu_n$  and  $\nu_{12}$  best fit  $\nu_{RL}$ . The fits of  $\nu_{12}$  to  $\nu_n$  for  $S/N \geq 1$  both have  $\chi^2 = 0.21$ , and  $\sim 0.25 - 0.5$  for full sets.

This suggests that differences in the derivation of power spectra from light curves can lead to non-negligible frequency differences in the frequencies.

16CygAB: frequency differences different power spectra: best fits to  $\nu_{RD}, \nu_{RL}$



## Uncertainties in mode fitting - mode heights


As stated above my fits to the power spectra had mode height ratios taken to be constant and with (my standard) values  $h_1/h_0 = 1.554$ ,  $h_2/h_0 = 0.582$ ,  $h_3/h_0 = 0.040$  which correspond to visibility coefficients for a limb darkening law  $f(\mu) = 0.3 + 0.7\mu$  (cf Roxburgh & Voontsov, 2006)<sup>1</sup>. I here explore the consequences of alternative fitting models on the results for 16CygB using Davies's power spectrum. I took the following models:

1. No  $\ell = 3$  modes: The standard values  $h_1/h_0 = 1.554$ ,  $h_2/h_0 = 0.582$  but setting  $h_3 = 0$
2. Highest ratios :  $h_1/h_0 = 1.688$ ,  $h_2/h_0 = 0.800$ ,  $h_3/h_0 = 0.109$  , limb darkening law  $f(\mu) = \mu$
3. Lowest ratios :  $h_1/h_0 = 1.333$ ,  $h_2/h_0 = 0.313$ ,  $h_3/h_0 = 0$  , limb darkening law  $f(\mu) = 1$ .
4. Free ratios :  $h_1/h_0$ ,  $h_2/h_0$ ,  $h_3/h_0$  unconstrained but lying between the values in 2) and 3) above. (*Davies et al (2015) take mode heights to be free parameters in their fits to 16CygA&B.*)

The results are displayed in the following diagram which gives the differences in frequencies relative to my standard fits  $\nu_{RD}$ . Neglecting the  $\ell = 3$  modes has a negligible effect on the frequencies but of course decreases the quality of fit of the model to the power spectrum Only the lowest ratios have a substantial effect on the values of the frequencies, the difference to the standard values decreasing as the height ratios are increased. Letting the ratios be free of course gives the best quality of fit to the power spectrum since there are more adjustable parameters - but I question whether this is reasonable. It may possibly be justified on the grounds that mode heights are not simply given by limb darkening and may vary with frequency, taking them as free parameters gives some idea of the uncertainties in the frequencies due to uncertainties in mode height ratios.

In my fitting routine I include  $\ell = 0, 2$  mode pairs at either end of the range to allow for the contribution of tails outside the frequencies to be determined - changing the constraints on these end values has negligible effect of the values of the frequencies of the modes whose values are sought,

16CygB: frequency differences between iwr fits to Davies' pspec  $\nu_{RD}$  (standard ratios)





## Uncertainties in rotation

The fits to Davies's power spectra for 16CygA&B ( $\nu_{RD}$ ) had rotation and inclination parameters taken as the central values reported in Davies et al (2015):  $(\omega, i) = (0.495, 56)$  for A, and  $(0.466, 36)$  for B, corresponding to  $\omega \sin i = 0.411$  for A and 0.274 for B (my standard values). Here I examine how the value of frequencies depends on the estimation of  $(\omega, i)$ . The figure below compares the frequencies of 3 fits to Davies's power spectrum for 16CygA to  $\nu_{RD}$ . All fits had my standard fixed mode height ratios corresponding to a limb darkening law  $f(\mu) = 0.3 + 0.7\mu$ . I give 3 examples:

- 1) pole on and/or no rotation;  $i=0$
- 2) equator on with standard  $\omega \sin i = 0.411$ ,  $i=90$
- 3) my best fit values

The top 2 panels in the figure show the frequency differences and  $\chi^2$  of the fits to my reference values  $\nu_{RD}$ . The pole on case is a poor fit ( $\chi^2 = 0.51$ ), the equator on case is better. The 3rd panel shows the difference between the  $i = 0$  and  $i = 90$  cases which gives some idea of the variation in frequencies with assumed inclination. A more detailed analysis gave  $\chi^2 = 0.5, 0.8, 0.6, 0.1$  for  $i = 10, 20, 30, 40$  all with the reference value of  $\omega \sin i = 0.411$ . The final panel compares the best fit values given by searching in a 2-dimensional mesh  $(\omega \sin i, i)$  to find the minimum. The values of  $(\omega, i) = (0.494, 52)$  are compatible with those of Davies in spite of the fact that  $\omega \sin i = 0.388$  is outside their estimated error bars ( $\omega \sin i = 0.411 \pm 0.013$ ). The difference in frequencies is negligible;  $\chi^2 = 0.004$ .

Davies et al took free mode height ratios so I repeated the analysis with free ratios; in this case the best fit had  $(\omega, i) = (0.508, 52)$ ,  $\omega \sin i = 0.400$ ,  $\chi^2 = 0.003$ , but somewhat larger differences at low  $i$

I did the same for 16CygB obtaining  $(\omega, i) = (0.339, 48)$ ,  $\omega \sin i = 0.252$ ,  $\chi^2 = 0.024$  with fixed height ratios; and  $(\omega, i) = (0.346, 50)$ ,  $\omega \sin i = 0.265$ ,  $\chi^2 = 0.048$  with free height ratios. Here the difference with Davies's values is larger but still the difference in splitting is very small. I note that my best fits for A&B have almost the same inclinations  $i = 50 \pm 2^\circ$ .

